



ACADEMIC
PRESS

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Advances in Mathematics 172 (2002) 151–157

ADVANCES IN
Mathematics

<http://www.elsevier.com/locate/aim>

A sharp bound for the Castelnuovo–Mumford regularity of subspace arrangements

Harm Derksen¹ and Jessica Sidman*

Department of Mathematics, University of Michigan, 525 East University Avenue, Ann Arbor, MI 48109-1109, USA

Received 2 January 2002; accepted 30 April 2002

Abstract

We show that the ideal of an arrangement of d linear subspaces of projective space is d -regular in the sense of Castelnuovo and Mumford, answering a question of B. Sturmfels. In particular, this implies that the ideal of an arrangement of d subspaces is generated in degrees less than or equal to d .

© 2002 Elsevier Science (USA). All rights reserved.

MSC: primary 13D02; secondary 52C35

0. Introduction

Over the past 20 years rapid advances in computational algebraic geometry have generated increasing amounts of interest in quantifying the “complexity” of ideals and modules. For a finitely generated module M over a polynomial ring $S = k[x_0, \dots, x_n]$ with k an arbitrary field, we say that M is r -regular (in the sense of Castelnuovo and Mumford) if the i th syzygy module of M is generated in degrees less than or equal to $r + i$. The notion of the regularity of M is key in determining the amount of computational resources that working with M requires. In this paper we will prove that the ideal of an arrangement of d linear spaces in \mathbb{P}^n is d -regular, answering a question posed by B. Sturmfels.

There is a mysterious gap in our understanding of the behavior of the regularity of ideals. Namely, it is known that in general the regularity of an ideal may be as bad as

*Corresponding author.

E-mail addresses: hderksen@math.lsa.umich.edu (H. Derksen), jsidman@umich.edu (J. Sidman).

¹Partially supported by NSF Grant DMS 0102193.

doubly exponential in the degrees of its minimal generators and the number of variables in the ambient ring, and this is essentially the worst case [4,12,13,16,20]. However, for an ideal I corresponding to a smooth variety in characteristic zero, the regularity of I is linear in terms of geometric data ([2,5,14,15,18]). From this perspective, the ideals of subspace arrangements provide a test case for the behavior of the regularity of ideals of reduced but possibly singular schemes.

Ideals of subspace arrangements also have connections with invariant theory and combinatorics. In [17], Noether gave a proof which showed that for the action of a finite group G on a polynomial ring $k[V]$ induced by a representation of G on V , the generators of $k[V]^G$ have degree less than or equal to the order of G as long as the characteristic of k is zero or if the characteristic is positive and greater than the order of G . For some time it was an open question as to whether or not the same result was true in positive characteristic with the stipulation that the characteristic of k not divide the order of G . Although the statement was recently shown to be true (see [8,10,11] for details), our main theorem yields an additional proof completing the circle of ideas introduced by the first author in [7]. Additionally, the ideals of subspace arrangements are studied by combinatorialists in relation to the computation of Betti numbers of the complement of an arrangement of linear subspaces in affine space [1].

The ideals of subspace arrangements have an appealing inductive structure and are algebraically closely related to the products of the linear ideals defining the constituent subspaces of the arrangement. The second author has used sheaf-theoretic techniques to study subspace arrangements of low dimension [19]. The proof given here is algebraic in nature and illustrates the close relationship of the intersection of linear ideals to their product, as it relies heavily on ideas introduced by Conca and Herzog to control the regularity of products of linear ideals in [6]. We recall basic notions related to regularity in Section 1 and present the proof of our result in Section 2.

1. Basic notions and conventions

Let $S = k[x_0, \dots, x_n]$ where k is an infinite field of arbitrary characteristic and $\mathfrak{m} = (x_0, \dots, x_n)$. Although we need to work over an infinite field, the regularity results also hold over a finite field k since passing to a field extension of k (which one may choose to be infinite) is exact. In this section we recall basic definitions and facts concerning regularity.

Of the various formulations of the definition of regularity, the following is the most illuminating from a computational point of view.

Definition 1.1. Let M be a finitely generated graded module over S . Take a minimal resolution of M by free graded S -modules:

$$0 \rightarrow \bigoplus_{\alpha=1}^{r_{n+1}} S(-e_{\alpha,n+1}) \rightarrow \cdots \rightarrow \bigoplus_{\alpha=1}^{r_0} S(-e_{\alpha,0}) \rightarrow M \rightarrow 0.$$

We say that M is r -regular if $\deg(e_{\alpha,i}) \leq r + i$ for all α, i and that the regularity of M , denoted $\text{reg}(M)$, is the least integer for which this holds.

One can also formulate the definition in terms of local cohomology and sheaf cohomology. The following theorem establishes the relationship between the regularity of an ideal and its associated sheaf.

Theorem 1.2 (Bayer and Mumford [2, Definition 3.2]). *Let I be a homogeneous ideal of S and let \tilde{I} be the associated coherent sheaf of ideals. Then the following conditions are equivalent:*

- (a) *The natural map $I_r \rightarrow H^0(\mathbb{P}^n, \tilde{I}(r))$ is an isomorphism and $H^i(\mathbb{P}^n, \tilde{I}(r - i)) = 0$, $1 \leq i \leq n$.*
- (b) *The natural maps $I_d \rightarrow H^0(\mathbb{P}^n, \tilde{I}(d))$ are isomorphisms for all $d \geq r$ and $H^i(\mathbb{P}^n, \tilde{I}(d)) = 0$, $d + i \geq r$, $i \geq 1$.*
- (c) *The ideal I is r -regular.*

A concept that arises frequently in the study of regularity is:

Definition 1.3. Let I be a homogeneous ideal of S . Then the saturation of I , denoted I^{sat} , is

$$I^{\text{sat}} := \{s \in S : \text{sm}^k \subseteq I, k \gg 0\}.$$

The saturation degree of I , denoted $\text{sat}(I)$, is the least integer k for which I agrees with I^{sat} in degrees k and higher.

It is also important to understand how regularity behaves in short exact sequences.

Lemma 1.4 (Eisenbud [9, Corollary 20.19]). *Let A , B , and C be finitely generated graded modules over S and let*

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

be a short exact sequence. Then

- (a) $\text{reg}(A) \leq \max\{\text{reg}(B), \text{reg}(C) + 1\}$.
- (b) $\text{reg}(B) \leq \max\{\text{reg}(A), \text{reg}(C)\}$.
- (c) $\text{reg}(C) \leq \max\{\text{reg}(A) - 1, \text{reg}(B)\}$.

The lemma below [3, Lemma 1.8] is the basis for a standard method of working inductively with regularity and is central to our proof in Section 2. We provide a proof here for the convenience of the reader. (Readers may also wish to consult [3] for a proof using local cohomology instead of sheaf cohomology.)

Lemma 1.5. *Let $I \subseteq k[x_0, \dots, x_n]$ be a homogeneous ideal, and suppose that x is a linear form that is a nonzerodivisor modulo I^{sat} . Then*

$$\text{reg}(I) = \max\{\text{reg}(I + (x)), \text{sat}(I)\}.$$

Proof. Note that the result will follow if we can show that I is r -regular if and only if I is saturated in degrees r and higher and $I + (x)$ is r -regular. It is crucial to observe here that $I \cap (x)$ can be rewritten as $(I : x)x$ and that if I is saturated in degrees r and higher then $(I : x)$ agrees with I in degrees r and higher.

For the forwards implication, assume that I is r -regular and note that since $\Gamma_*(\tilde{I}) = I^{\text{sat}}$, part (b) of Theorem 1.2 implies that an r -regular ideal has saturation degree less than or equal to r . To see that $I + (x)$ is r -regular, consider the short exact sequence

$$0 \rightarrow I \cap (x) \rightarrow I \oplus (x) \rightarrow I + (x) \rightarrow 0.$$

Since $(I : x)$ agrees with I in degrees r and higher, $(I : x)$ is r -regular which implies that $(I : x)x$ is $(r + 1)$ -regular. The r -regularity of $I + (x)$ follows from part (c) of Lemma 1.4.

For the reverse implication note that since $I + (x)$ is r -regular, its r th graded piece is isomorphic to $H^0(\mathbb{P}^n, (I + (x)) \sim (r))$. Furthermore, the r th graded piece of $I \oplus (x)$ injects into $H^0(\mathbb{P}^n, (I \oplus (x)) \sim (r))$ and surjects onto the r th graded piece of $I + (x)$. Therefore,

$$H^0(\mathbb{P}^n, (I \oplus (x)) \sim (r)) \rightarrow H^0(\mathbb{P}^n, (I + (x)) \sim (r))$$

is also a surjection. Together with the r -regularity of $I + (x)$ this implies that we have isomorphisms

$$H^i(\mathbb{P}^n, \tilde{I}(k - 1)) \rightarrow H^i(\mathbb{P}^n, \tilde{I}(k)) \oplus H^i(\mathbb{P}^n, (\tilde{x})(k))$$

for all $i > 0$ and $k \geq r - i + 1$. But now the vanishing of cohomology for $k \geq 0$ implies that $H^i(\mathbb{P}^n, \tilde{I}(r - i)) = 0$ for all $i > 0$. Since I is saturated in degrees r and higher by assumption, condition (a) of Theorem 1.2 is satisfied and I is r -regular. \square

2. The proof

We wish to prove the following theorem:

Theorem 2.1. *If X is an arrangement of d linear spaces in \mathbb{P}^n whose ideals are I_1, \dots, I_d then $I(X) = I_1 \cap \dots \cap I_d$ is d -regular.*

We should remark here that the statement of Theorem 2.1 is optimal. Indeed, one may take I to be the ideal of an arrangement of d lines that meet a fixed auxiliary line L in d distinct points. (L does not appear in the arrangement.) The hypersurface corresponding to any $f \in I$ must meet L in at least d points. Therefore, if f has degree strictly less than d , it must contain the line L . From this we can see that I must have at least one generator of degree at least d and that $\text{reg}(I) \geq d$.

Note that since $I_1 \cap \dots \cap I_d$ is saturated, our result would follow from Lemma 1.5 if we could show that $\text{reg}(I_1 \cap \dots \cap I_d + (x)) \leq d$. In fact, we prove:

Proposition 2.2. *Let I_1, \dots, I_d be the ideals of a linear arrangement in \mathbb{P}^n and let L be any linear ideal. Then*

$$I_1 \cap \dots \cap I_d + L$$

is d -regular.

Observe that $I_1 \cap \dots \cap I_d + \mathfrak{m} = \mathfrak{m}$ and is hence d -regular (in fact, 1-regular). This suggests that we ought to proceed by decreasing induction on $\text{ht}(L)$ (which is also the vector space dimension of the degree one part of L). The following lemma shows that bounding the saturation degree of $I_1 \cap \dots \cap I_d + L$ is actually the crucial computation:

Lemma 2.3. *If $I_1 \cap \dots \cap I_d + L'$ is d -regular for any linear ideal L' with $\text{ht}(L') > \text{ht}(L)$, then the d -regularity of $I_1 \cap \dots \cap I_d + L$ will follow if $I_1 \cap \dots \cap I_d + L$ is saturated in degrees d and higher.*

Proof. If $(I_1 \cap \dots \cap I_d + L)^{\text{sat}} = S$ then it is clear that if $I_1 \cap \dots \cap I_d + L$ is saturated in degrees d and higher then it is d -regular. If $(I_1 \cap \dots \cap I_d + L)^{\text{sat}} \neq S$ then choose a linear form x that is a nonzerodivisor modulo $(I_1 \cap \dots \cap I_d + L)^{\text{sat}}$. Using Lemma 1.5 we have

$$\text{reg}(I_1 \cap \dots \cap I_d + L) = \max\{\text{reg}(I_1 \cap \dots \cap I_d + L + (x)), \text{sat}(I_1 \cap \dots \cap I_d + L)\}.$$

Since x is a nonzerodivisor modulo $I_1 \cap \dots \cap I_d + L$, it cannot be contained in L . Therefore, $L + (x)$ is a linear ideal with height greater than $\text{ht}(L)$, so $I_1 \cap \dots \cap I_d + (L + (x))$ is d -regular by assumption. \square

Thus, the heart of the proof of Proposition 2.2 is a statement about saturation which we will control with the method used by Conca and Herzog [6].

Proof (Proposition 2.2). Set

$$J_i := I_1 \cap \dots \cap I_{i-1} \cap I_{i+1} \cap \dots \cap I_d + L$$

and

$$I := I_1 \cap \cdots \cap I_d + L.$$

Since $I_1 + L$ is 1-regular, we may assume inductively that J_i is $(d - 1)$ -regular.

Notice that if $\sum I_i + L \neq \mathfrak{m}$ then we can view the problem as living in a polynomial ring in fewer variables, say $S' = k[x_0, \dots, x_r]$ with $r < n$. In this case the generators of the ideal $I = I_1 \cap \cdots \cap I_d + L$ also live in S' . If we know that the ideal in S' is d -regular, then the d -regularity of the ideal in the larger ring follows by the flatness of $S'[x_{r+1}, \dots, x_n]$ over S' . But we may assume d -regularity in the smaller ring inductively since the result is trivial for \mathbb{P}^0 .

Finally, we show that I is saturated in degrees d and higher under the assumption that $\sum I_i + L = \mathfrak{m}$. Suppose there is a homogeneous form f of degree greater than or equal to d such that $f\mathfrak{m}^s \subseteq I$ for some integer $s > 0$. Choose a linear form x that is not in L and is a nonzerodivisor modulo all of the J_i^{sat} . Since $\text{ht}(L + (x)) > \text{ht}(L)$, by induction we see that $f \in I + (x)$ so $f = h + xf'$ with $h \in I$. Replace f by xf' , and note that $xf'\mathfrak{m}^s \subseteq I \subseteq J_i$ implies that $xf' \in J_i^{\text{sat}}$ for each i . Now, since x is a nonzerodivisor modulo J_i^{sat} , $xf' \in J_i^{\text{sat}}$ implies that $f' \in J_i^{\text{sat}}$. Additionally, we know that J_i is saturated in degrees $d - 1$ and higher which implies that $f' \in J_i$. Write $x = \sum x_i + x'$ where $x_i \in I_i$ and $x' \in L$. Observe that $x_i f' + x' f' \in I_i J_i \subseteq I$ for each i which implies that $xf' \in I$. Therefore, we have shown that I contains any element of degree d or higher in its saturation. The proposition now follows from Lemma 2.3. \square

Acknowledgments

We are very grateful to Bernd Sturmfels for suggesting the collaboration and Rob Lazarsfeld for his support, encouragement, and guidance. We are also deeply indebted to Aldo Conca and Jürgen Herzog for their many kind communications and in particular for sending us early versions of their preprints. Additionally, we thank Mel Hochster and Tamon Stephen for help searching the literature in commutative algebra and combinatorics, respectively, Dan Rogalski for helpful discussions, and Lawrence Ein and Karen Smith for their comments on the proof. The second author also thanks the NATO for its support of the *Workshop on Exterior Algebras and other new directions in Algebraic Geometry, Commutative Algebra, and Combinatorics* held in Sicily in September of 2001.

References

- [1] J. Alvarez Montaner, R. García López, S. Zarzuela Armengou, Local cohomology, arrangements of subspaces and monomial ideals, arXiv:math.AG0011181 v1 22 Nov 2000.
- [2] D. Bayer, D. Mumford, What Can Be Computed in Algebraic Geometry?, *Computational Algebraic Geometry and Commutative Algebra* (Cortona, 1991), Cambridge University Press, Cambridge, 1993, pp. 1–48.

- [3] D. Bayer, M. Stillman, A criterion for detecting m -regularity, *Invent. Math.* 87 (1987) 1–11.
- [4] D. Bayer, M. Stillman, On the complexity of computing syzygies, *J. Symbolic Comput.* 6 (2–3) (1988) 135–147 (Computational aspects of commutative algebra).
- [5] A. Bertram, L. Ein, R. Lazarsfeld, Vanishing theorems, a theorem of Severi, and the equations defining projective varieties, *J. Amer. Math. Soc.* 4 (3) (1991) 587–602.
- [6] A. Conca, J. Herzog, Castelnuovo–Mumford regularity of products of ideals, *Collectanea Mathematica*, in preparation.
- [7] H. Derksen, Computation of invariants for reductive groups, *Adv. Math.* 141 (2) (1999) 366–384.
- [8] H. Derksen, G. Kemper, *Computational Invariant Theory*, Book manuscript.
- [9] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Vol. 150, Springer, Berlin, 1995.
- [10] P. Fleischmann, The Noether bound in invariant theory of finite groups, *Adv. Math.* 156 (1) (2000) 23–32.
- [11] J. Fogarty, On Noether’s bound for polynomial invariants of a finite group, *Electron. Res. Announc. Amer. Math. Soc.* 7 (2001) 5–7 (electronic).
- [12] A. Galligo, Theoreme de division et stabilite en geometrie analytique locale, *Ann. Inst. Fourier (Grenoble)* 29 (1979) 107–184.
- [13] M. Giusti, Some effectivity problems in polynomial ideal theory, *EUROSAM 84* (Cambridge, 1984), Springer, Berlin, 1984, pp. 159–171.
- [14] L. Gruson, R. Lazarsfeld, C. Peskine, On a theorem of Castelnuovo, and the equations defining space curves, *Invent. Math.* 72 (3) (1983) 491–506.
- [15] R. Lazarsfeld, A sharp Castelnuovo bound for smooth surfaces, *Duke Math. J.* 55 (2) (1987) 423–429.
- [16] E.W. Mayr, A.R. Meyer, The complexity of the word problems for commutative semigroups and polynomial ideals, *Adv. in Math.* 46 (3) (1982) 305–329.
- [17] E. Noether, Der Endlichkeitssatz der Invarianten endlicher Gruppen. *Math. Ann.* (1916) 89–92.
- [18] H.C. Pinkham, A Castelnuovo bound for smooth surfaces, *Invent. Math.* 83 (2) (1986) 321–332.
- [19] J. Sidman, On the Castelnuovo–Mumford regularity of products of ideal sheaves, preprint.
- [20] C.-K. Yap, A new lower bound construction for commutative Thue systems with applications, *J. Symbolic Comput.* 12 (1) (1991) 1–27.