

Subject	Trial	Habituation	<i>Self-Feeding</i>	<i>Other-Feeding</i>
AH	1	313.07	np	np
	2	305.77	np	np
	3	222.00	np	np
	4	258.23	np	np
	5	--	np	np
	6	--	np	np
FL	1	382.33	np	3.10
	2	195.23	np	9.77
	3	182.53	np	8.03
	4	194.37	np	45.23
	5	--	np	np
	6	--	np	np
HG	1	9.97	np	np
	2	183.40	np	np
	3	15.80	np	np
	4	29.07	np	8.90
	5	--	np	5.70
	6	--	np	7.20
JM	1	182.43	np	14.47
	2	42.93	np	8.47
	3	16.27	np	4.90
	4	5.00	np	12.27
	5	--	np	7.90
	6	--	np	5.20
MD	1	180.83	np	np
	2	184.83	np	np
	3	25.27	np	np
	4	1.77	np	np
	5	--	np	np
	6	--	np	np
NN	1	216.67	np	10.83
	2	178.77	np	np
	3	186.43	np	np
	4	4.13	np	6.07
	5	--	np	np
	6	--	np	np

Table S.1. Latency (in seconds) to pull rope in Habituation and Pretest sessions by subject. Trials in which subjects did not pull the rope are denoted “np”.




Figure S.1. Side-view of the testing setup in (a) Self-Feeding, and (b) Other-Feeding trials in the Comprehension Pretest.