



Food craving as a mediator between addictive-like eating and problematic eating outcomes



Michelle A. Joyner^{a,*}, Ashley N. Gearhardt^a, Marney A. White^{b,c}

^a Department of Psychology, University of Michigan, 530 Church Street, Ann Arbor, MI 48109-1043, USA

^b Department of Psychiatry, Yale School of Medicine, 300 George Street, New Haven, CT 06511, USA

^c Department of Chronic Disease Epidemiology, Yale School of Public Health, 60 College Street, New Haven, CT 06511, USA

ARTICLE INFO

Article history:

Received 23 April 2015

Received in revised form 25 June 2015

Accepted 29 July 2015

Available online 4 August 2015

Keywords:

Craving
Food addiction
Binge eating
BMI

ABSTRACT

Background: There is growing interest and debate about whether an addictive process contributes to problematic eating outcomes, such as obesity. Craving is a core component of addiction, but there has been little research on the relationship between addictive-like eating, craving, and eating-related concerns. In the current study, we examine the effect of both overall food craving and craving for different types of food on the relationship between addictive-like eating symptoms and elevated body mass index (BMI) and binge eating episodes.

Methods: In a community sample ($n = 283$), we conducted analyses to examine whether overall craving mediated the association between addictive-like eating and elevated BMI, as well as binge eating frequency. We also ran separate mediational models examining the indirect effect of cravings for sweets, fats, carbohydrates, and fast food fats on these same associations.

Results: Overall food craving was a significant partial mediator in the relationships between addictive-like eating and both elevated BMI and binge eating episodes. Cravings for sweets and other carbohydrates significantly mediated the relationship between addictive-like eating and binge eating episodes, while cravings for fats significantly mediated the relationship between addictive-like eating and elevated BMI.

Conclusions: Craving appears to be an important component in the pathway between addictive-like eating and problematic eating outcomes. The current results highlight the importance of further evaluating the role of an addictive process in problematic eating behaviors and potentially targeting food cravings in intervention approaches.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

The theory that certain foods can trigger an addictive process has recently gained attention (Avena, Rada, & Hoebel, 2008; Davis & Carter, 2009; Gearhardt, Corbin, & Brownell, 2009; Gold, Graham, Cocores, & Nixon, 2009; Ifland et al., 2009; Pelchat, 2002), although it remains controversial (Avena, Gearhardt, Gold, Wang, & Potenza, 2012; Ziauddeen, Farooqi, & Fletcher, 2012; Ziauddeen & Fletcher, 2013). Food and drug rewards appear to act on similar neural pathways (Berridge, Ho, Richard, & DiFeliceantonio, 2010; Volkow, Wang, Fowler, & Telang, 2008). Rats exhibit behaviors resembling symptoms of addiction in response to sugar consumption (Avena et al., 2008; Colantuoni et al., 2001; Rada, Avena, & Hoebel, 2005). Craving appears to play an important role in compulsive use across addictive substances, predicting negative outcomes such as increased use and earlier relapse in alcohol, cocaine, and tobacco use (Bottlender & Soyka, 2004; Donny, Griffin, Shiffman, & Sayette, 2008; Epstein, Marrone, Heishman, Schmittner, &

Preston, 2010; Flannery, Poole, Gallop, & Volpicelli, 2003; Oslin, Cary, Slaymaker, Colleran, & Blow, 2009; Weiss et al., 2003). Examining the association of addictive-like eating and food craving is important in evaluating the “food addiction” hypothesis. The Yale Food Addiction Scale (YFAS) was developed to assess addictive-like eating in humans by applying *DSM-IV-TR* substance dependence criteria to the consumption of certain foods (Gearhardt et al., 2009). YFAS “food addiction” symptoms are related to elevated craving (Gearhardt, Rizk, & Treat, 2014; Meule & Kübler, 2012), as well as elevated BMI and more frequent binge eating episodes (Davis et al., 2011; Gearhardt, Boswell, & White, 2014). Food craving appears to be associated with eating related problems, such as binge eating and elevated body mass index (BMI) (Greeno, Wing, & Shiffman, 2000; Waters, Hill, & Waller, 2001; White & Grilo, 2005). However, fat and sugar cravings appear to have different associations with “food addiction” (Gearhardt, Rizk, & Treat, 2014) and obesity (Drewnowski, Kurth, Holden-Wiltse, & Saari, 1992).

To our knowledge, no prior studies have investigated whether higher food craving may be a pathway through which addictive-like eating is associated with eating-related problems and whether this differs by the type of food that is craved. In the current study, we aim to test food craving as a mediator between addictive-like eating and both

* Corresponding author at: Department of Psychology, University of Michigan, 2257 East Hall, Ann Arbor, MI 48109-1043. Tel.: +1 734 647 3873.
E-mail address: majoy@umich.edu (M.A. Joyner).

BMI and binge eating episodes. Additionally, we examine whether cravings for different types of foods (sweets, carbohydrates, fats, and fast food fats) may differentially mediate these associations.

2. Methods

2.1. Participants

The study was approved by the Yale Institutional Review Board. Participants ($n = 283$) were drawn from a sample of 420 community volunteers and were included in this study if they provided complete data on key study measures. Participants were on average 34.9 years old (range 18–69). The sample was 15.9% male ($n = 45$) and 83.0% female ($n = 235$), and 3 participants did not report gender. The racial/ethnic distribution for the study sample was: 73.1% Caucasian, 9.5% Hispanic, 7.8% African American, 4.9% Asian, and 3.9% “other.” Two participants did not report race/ethnicity. The participants' body weight ranged from underweight to severely obese (BMI range 14.75 to 66.07) with the average BMI in the overweight category ($M = 28.44$, $SD = 9.00$).

2.2. Assessments and measures

Participants provided basic demographic information and completed a battery of self-report measures. Self-reported height and weight were used to compute participant BMI (kg/m^2).

The *Yale Food Addiction Scale* (Gearhardt et al., 2009) measures signs of “addiction” toward certain types of food (e.g. high in fat, high in sugar) based on the criteria for substance dependence as stated in the *DSM-IV-TR* (American Psychiatric Association, 2000). The YFAS “symptom count” score reflects the number of addiction-like criteria (ranging 0–7) endorsed. In the current sample, the YFAS exhibited adequate internal consistency ($\alpha = .77$). The mean YFAS symptom count score in the current sample was 3.02 ($SD = 2.00$).

The *Food Craving Inventory* (FCI) (White, Whisenhunt, Williamson, Greenway, & Netemeyer, 2002) is a 28-item measure that assesses the frequency of cravings for specific foods. The FCI yields a score of total food craving, as well as four subscales measuring cravings for high fats (e.g., bacon), carbohydrates/starches (e.g., bread), sweets (e.g., cookies), and fast food fats (e.g., hamburgers). Total and subscale scores each have a possible range of 1–5. In the current sample, the FCI exhibited excellent internal consistency ($\alpha = .95$). The mean total craving score in the current sample was 2.09 ($SD = .83$). Mean subscale scores were as follows: fats ($M = 1.75$, $SD = .88$), fast food fats ($M = 2.36$, $SD = .99$), sweets ($M = 2.32$, $SD = .99$), and carbohydrates ($M = 1.91$, $SD = .90$).

The *Eating Disorder Examination Questionnaire* (EDE-Q), a well-established measure of eating disorder psychopathology (Fairburn & Beglin, 1994; Luce & Crowther, 1999), was used to assess the number of binge eating episodes. The item used asked participants on how many times in the past 28 days they had experienced a binge eating episode (i.e., eaten an unusually large amount of food and felt a loss of control while doing so). In the current sample, the number of binge eating episodes reported ranged from 0–28, with a mean of 2.60 ($SD = 5.18$).

2.3. Statistical analyses

Binge eating episodes were skewed, thus we removed four outliers ($SD > 2$). Age was significantly correlated with BMI ($r = .303$, $p < .001$) and was included as a covariate in all analyses.

To test the hypothesized mediational models (e.g. addictive-like eating symptoms \rightarrow food craving \rightarrow BMI), we followed the guidelines described by Baron and Kenny (1986). To further examine any mediational effects, we employed the bootstrapping method with 10,000 samples described by Preacher and Hayes (2008). We used the completely standardized indirect effect (ab_{cs}) described by Preacher

and Kelley (2011) to compare the effect sizes of statistically significant indirect effects. Effect sizes can be interpreted as small (.01), medium (.09) or large (.25) (Kenny, 2014).

3. Results

3.1. Addictive-like eating and BMI

Addictive-like eating was significantly associated with both BMI and food craving (i.e., FCI–Total score), and food craving was significantly associated with BMI (see Fig. 1). Food craving was a significant partial mediator between addictive-like eating and BMI ($B = .46$, $SE = .24$, $95\% \text{ CI} = .03-.99$, $ab_{cs} = .11$).

3.2. Addictive-like eating and binge eating episodes

Food craving was a significant partial mediator between addictive-like eating and binge eating episodes ($B = .29$, $SE = .14$, $95\% \text{ CI} = .06-.62$, ($ab_{cs} = .12$) (see Fig. 2).

3.3. Food cravings for specific food types

3.3.1. Fats

Fats craving was a significant partial mediator between addictive-like eating behaviors and BMI illustrated through both regression analysis ($\beta = .155$, $p = .018$, $\text{adj. } R^2 = .215$) and bootstrapping ($B = .31$, $SE = .19$, $95\% \text{ CI} = .01-.75$, $ab_{cs} = .08$).

Craving for fats did not significantly predict the number of binge eating episodes when addictive-like eating was included in the model ($\beta = .108$, $p = .067$, $\text{adj. } R^2 = .356$) and did not significantly mediate the relationship according to bootstrapping ($B = .18$, $SE = .10$, $95\% \text{ CI} = -.04 \text{ to } .34$, $ab_{cs} = .05$).

3.3.2. Fast food fats

Craving for fast food fats did not significantly predict BMI when addictive-like eating was included ($\beta = .098$, $p = .135$, $\text{adj. } R^2 = .203$) and did not significantly mediate this relationship according to bootstrapping ($B = .21$, $SE = .14$, $95\% \text{ CI} = -.04 \text{ to } .52$, $ab_{cs} = .05$). Craving for fast food fats also did not significantly predict the number of binge eating episodes when addictive-like eating was included in the model ($\beta = .097$, $p = .095$, $\text{adj. } R^2 = .354$), and did not significantly mediate this relationship according to bootstrapping ($B = .11$, $SE = .07$, $95\% \text{ CI} = -.01 \text{ to } .27$, $ab_{cs} = .04$).

3.3.3. Sweets

Craving for sweets did not significantly predict BMI when addictive-like eating was included in the model ($\beta = .134$, $p = .067$, $\text{adj. } R^2 = .207$) and did not significantly mediate this relationship according to bootstrapping ($B = .35$, $SE = .20$, $95\% \text{ CI} = -.04 \text{ to } .77$, $ab_{cs} = .09$).

Sweets craving was a significant partial mediator between addictive-like eating behaviors and number of binge eating episodes illustrated




Fig. 1. Standardized regression coefficients for the relationship between addictive-like eating and BMI as mediated by food craving. The standardized regression coefficient between addictive-like eating and BMI controlling for food craving is in parentheses. * $p < .05$, ** $p < .01$.




Fig. 2. Standardized regression coefficients for the relationship between addictive-like eating and binge eating episodes as mediated by food craving. The standardized regression coefficient between addictive-like eating and binge eating episodes controlling for food craving is in parentheses. * $p < .05$, ** $p < .01$.

through both regression analysis ($\beta = .206$, $p = .001$, adj. $R^2 = .374$) and bootstrapping ($B = .28$, $SE = .12$, 95% $CI = .08-.56$, $ab_{cs} = .12$).

3.3.4. Carbohydrates

Craving for carbohydrates did not significantly predict BMI when addictive-like eating was included in the model ($\beta = .130$, $p = .060$, adj. $R^2 = .212$) and did not significantly mediate this relationship according to bootstrapping ($B = .31$, $SE = .21$, 95% $CI = -.06$ to $.78$, $ab_{cs} = .08$).

Carbohydrates craving was a significant partial mediator between addictive-like eating behaviors and number of binge eating episodes illustrated through both regression analysis ($\beta = .178$, $p = .003$, adj. $R^2 = .369$) and bootstrapping ($B = .22$, $SE = .12$, 95% $CI = .03-.52$, $ab_{cs} = .09$).

4. Discussion

In the current study, food craving was found to partially mediate the relationships between addictive-like eating and BMI and between addictive-like eating and number of binge eating episodes. Thus, consistent with an addiction perspective, craving appears to be an important component in the pathway between addictive-like eating and eating-related problems. Cravings for different types of foods differentially mediated the association between addictive-like eating and eating-related concerns. While many palatable foods contain high amounts of sugar and fat, these findings suggest that there may be differences based on the prominence of each component in a particular food. Cravings for fats mediated the relationship between addictive-like eating and BMI, but not binge eating episodes. Fat consumption has been more closely related to weight gain and somatosensory processing than sugar, but may contribute less to behavioral outcomes related to addiction, such as bingeing (Avena, Rada, & Hoebel, 2009; Stice, Burger, & Yokum, 2013). Cravings for fast food fats did not significantly mediate the relationships between addictive-like eating and either BMI or binge eating episodes, possibly because this category of foods is too specific. Cravings for sweets significantly mediated the relationship between addictive-like eating and binge eating episodes, but not BMI. Sugar has been implicated in activating reward-related neural pathways and is related to the development of addictive-like eating behaviors, but not weight gain in rats (Avena et al., 2008; Stice et al., 2013). Cravings for foods high in non-sugar carbohydrates also significantly mediated the relationship between food addiction and binge eating episodes, but not BMI. Starchy carbohydrates are broken down into sugars during digestion and increase blood sugar levels (Ludwig, 2002), so it is possible that refined carbohydrates could trigger an addictive process much like sugar. Therefore, cravings for sugar or non-sugar carbohydrates appear to be more closely associated with addictive-like eating and bingeing, while cravings for fat appear to be more closely related to elevated BMI.

This study does have some limitations. Due to the cross-sectional design, we cannot make causal attributions. Future research should employ a longitudinal design. This sample consisted of volunteers

recruited from an advertisement for a study on eating behavior and health, which could have led to some selection bias. The current sample also included mostly women, who tend to report higher rates of loss of control and binge eating (Striegel-Moore et al., 2009). These issues can be addressed in the future by replicating this study using a random and representative sample. BMI was calculated using self-reported height and weight and future studies would benefit from direct measurement of height and weight.

4.1. Conclusions

Akin to substances such as alcohol or cocaine (Bottlender & Soyka, 2004; Weiss et al., 2003), food craving appears to be associated with overconsumption in those experiencing addictive-like eating symptoms. Cravings for sweets and other high carbohydrate foods may be more closely associated with addictive-like eating behaviors such as bingeing, while cravings for fats appear to be more related to elevated BMI. Craving may be a particularly important intervention target for individuals who report greater addictive-like eating.

Role of funding sources

Partial funding for this study was provided by NIDDK Grant K23 DK071646. NIDDK had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

Contributors

Marney White designed and supervised the data collection and wrote the protocol. Michelle Joyner and Ashley Gearhardt developed the current study aims and conducted the statistical analyses. Michelle Joyner wrote the first draft of the manuscript and Ashley Gearhardt and Marney White aided in the revision of the manuscript. All authors contributed to and have approved the final manuscript.

Conflict of interest

All authors declare that they have no conflicts of interest.

References

- American Psychiatric Association (2000). *Diagnostic and statistical manual of mental disorders* (4th ed. text revision) (Washington DC).
- Avena, N. M., Gearhardt, A. N., Gold, M. S., Wang, G. J., & Potenza, M. N. (2012). Tossing the baby out with the bathwater after a brief rinse? The potential downside of dismissing food addiction based on limited data. *Nature Review Neuroscience*, 13(7), 514. <http://dx.doi.org/10.1038/nrn3212-c1> (author reply 514).
- Avena, N. M., Rada, P., & Hoebel, B. G. (2008). Evidence for sugar addiction: Behavioral and neurochemical effects of intermittent, excessive sugar intake. *Neuroscience & Biobehavioral Reviews*, 32(1), 20–39.
- Avena, N. M., Rada, P., & Hoebel, B. G. (2009). Sugar and fat bingeing have notable differences in addictive-like behavior. *Journal of Nutrition*, 139(3), 623–628. <http://dx.doi.org/10.3945/jn.108.097584>.
- Baron, R. M., & Kenny, D. A. (1986). The moderator–mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, 51(6), 1173–1182.
- Berridge, K. C., Ho, C. Y., Richard, J. M., & DiFeliceantonio, A. G. (2010). The tempted brain eats: Pleasure and desire circuits in obesity and eating disorders. *Brain Research*, 1350, 43–64. <http://dx.doi.org/10.1016/j.brainres.2010.04.003>.
- Bottlender, M., & Soyka, M. (2004). Impact of craving on alcohol relapse during, and 12 months following, outpatient treatment. *Alcohol and Alcoholism*, 39(4), 357–361. <http://dx.doi.org/10.1093/alcalc/agh073>.
- Colantuoni, C., Schwenker, J., McCarthy, J., Ladenheim, B., Cadet, J.-L., Schwartz, G. J., et al. (2001). Excessive sugar intake alters binding to dopamine and mu-opioid receptors in the brain. *NeuroReport*, 12(16), 3549–3552.
- Davis, C., & Carter, J. C. (2009). Compulsive overeating as an addiction disorder. A review of theory and evidence. *Appetite*, 53(1), 1–8. <http://dx.doi.org/10.1016/j.appet.2009.05.018>.
- Davis, C., Curtis, C., Levitan, R. D., Carter, J. C., Kaplan, A. S., & Kennedy, J. L. (2011). Evidence that 'food addiction' is a valid phenotype of obesity. *Appetite*, 57(3), 711–717. <http://dx.doi.org/10.1016/j.appet.2011.08.017>.
- Donny, E. C., Griffin, K. M., Shiffman, S., & Sayette, M. A. (2008). The relationship between cigarette use, nicotine dependence, and craving in laboratory volunteers. *Nicotine & Tobacco Research*, 10(3), 447–455. <http://dx.doi.org/10.1080/14622200801901906>.
- Drewnowski, A., Kurth, C., Holden-Wiltse, J., & Saari, J. (1992). Food preferences in human obesity: Carbohydrates versus fats. *Appetite*, 18, 207–221.
- Epstein, D. H., Marrone, G. F., Heishman, S. J., Schmittner, J., & Preston, K. L. (2010). Tobacco, cocaine, and heroin: Craving and use during daily life. *Addictive Behaviors*, 35(4), 318–324. <http://dx.doi.org/10.1016/j.addbeh.2009.11.003>.
- Fairburn, C. G., & Beglin, S. J. (1994). Assessment of eating disorders: Interview or self-report questionnaire? *International Journal of Eating Disorders*, 16(4), 363–370.

- Flannery, B. A., Poole, S. A., Gallop, R. J., & Volpicelli, J. R. (2003). Alcohol craving predicts drinking during treatment: An analysis of three assessment instruments. *Journal of Studies on Alcohol*, 64, 120–126.
- Gearhardt, A. N., Boswell, R. G., & White, M. A. (2014). The association of “food addiction” with disordered eating and body mass index. *Eating Behaviors*, 15(3), 427–433. <http://dx.doi.org/10.1016/j.eatbeh.2014.05.001>.
- Gearhardt, A. N., Corbin, W. R., & Brownell, K. D. (2009). Preliminary validation of the Yale Food Addiction Scale. *Appetite*, 52(2), 430–436. <http://dx.doi.org/10.1016/j.appet.2008.12.003>.
- Gearhardt, A. N., Rizk, M. T., & Treat, T. A. (2014). The association of food characteristics and individual differences with ratings of craving and liking. *Appetite*, 79, 166–173. <http://dx.doi.org/10.1016/j.appet.2014.04.013>.
- Gold, M. S., Graham, N. A., Cocores, J. A., & Nixon, S. J. (2009). Food addiction? *Journal of Addiction Medicine*, 3(1), 42–45.
- Greeno, C. G., Wing, R. R., & Shiffman, S. (2000). Binge antecedents in obese women with and without binge eating disorder. *Journal of Consulting and Clinical Psychology*, 68(1), 95–102. <http://dx.doi.org/10.1037//0022-006x.68.1.95>.
- Ifland, J. R., Preuss, H. G., Marcus, M. T., Rourke, K. M., Taylor, W. C., Burau, K., et al. (2009). Refined food addiction: A classic substance use disorder. *Medical Hypotheses*, 72(5), 518–526. <http://dx.doi.org/10.1016/j.mehy.2008.11.035>.
- Kenny, D. A. (2014). Mediation. from <http://davidakenny.net/cm/mediate.htm>
- Luce, K. H., & Crowther, J. H. (1999). The reliability of the Eating Disorder Examination–Self-report questionnaire version (EDE-Q). *International Journal of Eating Disorders*, 25, 349–351.
- Ludwig, D. S. (2002). The glycemic index: Physiological mechanisms relating to obesity, diabetes, and cardiovascular disease. *JAMA*, 287(18), 2414–2423.
- Meule, A., & Kübler, A. (2012). Food cravings in food addiction: The distinct role of positive reinforcement. *Eating Behaviors*, 13(3), 252–255. <http://dx.doi.org/10.1016/j.eatbeh.2012.02.001>.
- Oslin, D. W., Cary, M., Slaymaker, V., Collieran, C., & Blow, F. C. (2009). Daily ratings measures of alcohol craving during an inpatient stay define subtypes of alcohol addiction that predict subsequent risk for resumption of drinking. *Drug and Alcohol Dependence*, 103(3), 131–136. <http://dx.doi.org/10.1016/j.drugalcdep.2009.03.009>.
- Pelchat, M. L. (2002). Of human bondage: Food craving, obsession, compulsion, and addiction. *Physiology & Behavior*, 76, 347–352.
- Preacher, K. J., & Hayes, A. F. (2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods*, 40(3), 879–891. <http://dx.doi.org/10.3758/brm.40.3.879>.
- Preacher, K. J., & Kelley, K. (2011). Effect size measures for mediation models: Quantitative strategies for communicating indirect effects. *Psychological Methods*, 16(2), 93–115. <http://dx.doi.org/10.1037/a0022658.supp>.
- Rada, P., Avena, N. M., & Hoebel, B. G. (2005). Daily bingeing on sugar repeatedly releases dopamine in the accumbens shell. *Neuroscience*, 134(3), 737–744. <http://dx.doi.org/10.1016/j.neuroscience.2005.04.043>.
- Stice, E., Burger, K. S., & Yokum, S. (2013). Relative ability of fat and sugar tastes to activate reward, gustatory, and somatosensory regions. *American Journal of Clinical Nutrition*, 98(6), 1377–1384. <http://dx.doi.org/10.3945/ajcn.113.069443>.
- Striegel-Moore, R. H., Rosselli, F., Perrin, N., DeBar, L., Wilson, G. T., May, A., et al. (2009). Gender difference in the prevalence of eating disorder symptoms. *International Journal of Eating Disorders*, 42(5), 471–474. <http://dx.doi.org/10.1002/eat.20625>.
- Volkow, N. D., Wang, G. J., Fowler, J. S., & Telang, F. (2008). Overlapping neuronal circuits in addiction and obesity: Evidence of systems pathology. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 363(1507), 3191–3200. <http://dx.doi.org/10.1098/rstb.2008.0107>.
- Waters, A., Hill, A., & Waller, G. (2001). Internal and external antecedents of binge eating episodes in a group of women with bulimia nervosa. *International Journal of Eating Disorders*, 29, 17–22.
- Weiss, R. D., Griffin, M. L., Mazurick, C., Berkman, B., Gastfriend, D. R., Frank, A., et al. (2003). The relationship between cocaine craving, psychosocial treatment, and subsequent cocaine use. *The American Journal of Psychiatry*, 160(7), 1320–1325.
- White, M. A., & Grilo, C. M. (2005). Psychometric properties of the Food Craving Inventory among obese patients with binge eating disorder. *Eating Behaviors*, 6(3), 239–245. <http://dx.doi.org/10.1016/j.eatbeh.2005.01.001>.
- White, M. A., Whisenhunt, B. L., Williamson, D. A., Greenway, F. L., & Netemeyer, R. G. (2002). Development and validation of the Food-Craving Inventory. *Obesity Research*, 10(2), 107–114.
- Ziauddeen, H., Farooqi, I. S., & Fletcher, P. C. (2012). Obesity and the brain: How convincing is the addiction model? *Nature Review Neuroscience*, 13(4), 279–286. <http://dx.doi.org/10.1038/nrn3212>.
- Ziauddeen, H., & Fletcher, P. C. (2013). Is food addiction a valid and useful concept? *Obesity Reviews*, 14(1), 19–28. <http://dx.doi.org/10.1111/j.1467-789X.2012.01046.x>.